Simple Is Easy

Discover why oral BAFIERTAM[®] is the safe, effective, and simple choice for treating your relapsing multiple sclerosis.



Indication and Select Important Safety Information

What is BAFIERTAM (monomethyl fumarate)?

- BAFIERTAM is a prescription medicine used to treat relapsing forms of multiple sclerosis (MS), to include clinically isolated syndrome, relapsing-remitting disease, and active secondary progressive disease, in adults.
- It is not known if BAFIERTAM is safe and effective in children.

Who should not take BAFIERTAM?

- Do not take BAFIERTAM if you: have had an allergic reaction (such as welts, hives, swelling of the face, lips, mouth or tongue, or difficulty breathing) to monomethyl fumarate, dimethyl fumarate, diroximel fumarate, or any of the ingredients in BAFIERTAM.
- Do not take BAFIERTAM if you are taking dimethyl fumarate or diroximel fumarate.

Please see additional Important Safety Information on pages 8-10 and the accompanying full <u>Prescribing Information</u> and <u>Patient Information</u>.

What Makes BAFIERTAM Different?

BAFIERTAM is one of a proven class of oral drugs called fumarates, which have long been known for their **safety and effectiveness**. BAFIERTAM is different and easier to take:

- Can be taken with or without food
- No need to count calories or restrict or change your eating or drinking habits
- May be taken without dose adjustments or restrictions in people with kidney problems
- Leaves the bloodstream within 24 hours, so easy to start and stop as needed

Small soft-gel capsules make BAFIERTAM easier to take



BAFIERTAM (MMF) 190 mg twice daily¹



Pills are shown to scale but not actual size

Other fumarates (DMF/DRF) 240 mg for DMF and 462 mg for DRF twice daily^{2,3}


1. BAFIERTAM. Prescribing Information. Banner Life Sciences LLC; 2021. **2.** TECFIDERA. Prescribing Information. Biogen Inc; 2022. **3.** VUMERITY. Prescribing Information. Biogen Inc; 2022. **4.** Gold R, Arnold DL, Bar-Or A, et al. Long-term safety and efficacy of dimethyl fumarate for up to 13 years in patients with relapsing-remitting multiple sclerosis: Final ENDORSE study results. *Multiple Sclerosis Journal.* 2022;28(5):801-816. doi:10.1177/13524585211037909



Learn More About the BAFIERTAM Difference BAFIERTAM.com/why-bafiertam

An Easy, Effective Way to Slow the Progression of Multiple Sclerosis (MS)

BAFIERTAM works to slow the overall disease progression of MS, while reducing the severity and frequency of symptoms. Based on a long-term study of DMF, patients experienced the equivalent of less than 2 relapses in 20 years⁴. The active ingredient (MMF) in BAFIERTAM was proven to reduce*:



Common Side Effects

BAFIERTAM shares the same side effects as other fumarates. Flushing and stomach upset are the most common, especially at the start of treatment, and usually decrease over time. Call your doctor if any of these symptoms bother you or do not go away.

Tip: Ask your doctor if taking a nonenteric-coated aspirin (up to 325 mg) 30 minutes before BAFIERTAM is safe for you. This has been shown to help limit flushing.



Tip: Taking BAFIERTAM without food may also relieve stomach upset.

*Based on studies conducted with DMF by Biogen Inc.

Please see additional Important Safety Information on pages 8-10 and the accompanying full <u>Prescribing Information</u> and <u>Patient Information</u>.



Is BAFIERTAM Right for You?

BAFIERTAM is in the most widely prescribed class of oral MS therapies. Whether you're starting your MS treatment or switching from a similar medication, BAFIERTAM may be an easy choice if you:


- Prefer an oral medication over an injection
- Have safety concerns regarding any existing health conditions, like kidney disease, or medications you may be taking
- Want the flexibility that comes with no dietary restrictions
- Are planning to or may become pregnant

Before you start, you'll have a simple blood test. It will be repeated about every 6 months to monitor your condition.

Twice-Daily Dosing

Remembering to take BAFIERTAM is as simple as remembering to eat breakfast and dinner or brushing your teeth each morning and night.

BAFIERTAM small gel capsules make dosing convenient and easy to fit into your lifestyle. Your doctor may prescribe a 7-day starter dose of 1 pill twice daily. Then, maintenance dosing is 2 pills twice daily.







Get All the Details About BAFIERTAM BAFIERTAM.com

Ask Your Doctor About BAFIERTAM

It's always best to discuss your specific condition and needs with your doctor before starting a new treatment. **These questions can help you start the conversation:**

- □ How does BAFIERTAM work?
- Can you explain how BAFIERTAM can delay the progression of physical disability?
- □ What impact will BAFIERTAM have on my relapses and MRI lesions?
- How is BAFIERTAM different from other fumarates?
- □ What side effects should I expect with BAFIERTAM?
- □ What can I do to limit the onset of any side effects?
- Do I need to be concerned about what or when I eat when I take BAFIERTAM?
- □ Will BAFIERTAM interact with other prescription drugs I take?
- □ Is BAFIERTAM easy to swallow?
- □ How soon can I get pregnant if I discontinue BAFIERTAM?
- How easy is it to get started on BAFIERTAM?
- □ Is there specific medical testing needed to get started?
- □ What patient support is available with BAFIERTAM?

Use this guide to discuss BAFIERTAM with your doctor.

Please see additional Important Safety Information on pages 8-10 and the accompanying full <u>Prescribing Information</u> and <u>Patient Information</u>.



You're Not Alone: Banner Life Sciences® Is Here for You

When your doctor prescribes BAFIERTAM, Banner Patient Support is here to help throughout your time on treatment with:

- One-to-one support from a BAFIERTAM Patient Support Care Manager
- Assistance starting BAFIERTAM while your insurance benefits are confirmed
- Coordinating your insurance benefits with your specialty pharmacy
- Assessing and obtaining available financial assistance

Financial and Insurance Support for Eligible Patients



QuickStart Program Provides a 30-day supply of BAFIERTAM to start treatment if your insurance benefits are delayed.



Bridge Support Program* Helps you keep receiving BAFIERTAM if you experience a gap in commercial insurance coverage.



\$0 Copay Program[†] If you have commercial health insurance, you may be eligible to receive BAFIERTAM for a \$0 copay.



Visit our website for more information about BAFIERTAM Patient Support Programs. copay.BAFIERTAM.com

*Banner Life Sciences reserves the right to modify this program at any time without notice. Other limitations may apply

¹The \$0 Copay Program is for patients taking BAFIERTAM and is subject to an annual cap on the amount of assistance patients can receive. This offer is invalid for patients covered by any governmental program, including, without limitation, Medicaid, Medicare, VA, or TRICARE. Federal and state laws and other factors may prevent or otherwise restrict eligibility.



Your Next Step

Banner respects your privacy. To get started, please provide your consent to enroll in patient services by:

- Signing the enrollment form your doctor may use to prescribe BAFIERTAM, or
- Visiting our website, entering the Consent Portal, completing and signing the consent form, or



Visit allcareconsent.com

Calling us at 1-855-3BANNER (1-855-322-6637)



Once your doctor has prescribed BAFIERTAM, a Banner care manager will reach out to help you get started.

Please see additional Important Safety Information on pages 8-10 and the accompanying full <u>Prescribing Information</u> and <u>Patient Information</u>.



Indication and Important Safety Information

What is BAFIERTAM[®] (monomethyl fumarate)?

- BAFIERTAM is a prescription medicine used to treat relapsing forms of multiple sclerosis (MS), to include clinically isolated syndrome, relapsing-remitting disease, and active secondary progressive disease, in adults.
- It is not known if BAFIERTAM is safe and effective in children.

Who should not take BAFIERTAM?

- Do not take BAFIERTAM if you: have had an allergic reaction (such as welts, hives, swelling of the face, lips, mouth or tongue, or difficulty breathing) to monomethyl fumarate, dimethyl fumarate, diroximel fumarate, or any of the ingredients in BAFIERTAM.
- Do not take BAFIERTAM if you are taking dimethyl fumarate or diroximel fumarate.

Before taking and while you take BAFIERTAM, tell your doctor about all of your medical conditions, including if you:

- have liver problems
- have or have had low white blood cell counts or an infection
- are pregnant or plan to become pregnant. It is not known if BAFIERTAM will harm your unborn baby
- are breastfeeding or plan to breastfeed. It is not known if BAFIERTAM passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby while using BAFIERTAM.

Tell your doctor about all the medicines you take including prescription and over-thecounter medicines, vitamins, and herbal supplements.

What are the possible side effects of BAFIERTAM?

BAFIERTAM may cause serious side effects, including:

• allergic reaction (such as welts, hives, swelling of the face, lips, mouth or tongue, or difficulty breathing). Stop taking BAFIERTAM and get emergency medical help right away if you get any of these symptoms.

- **PML (progressive multifocal leukoencephalopathy)** a rare brain infection that usually leads to death or severe disability over a period of weeks or months.
 - Tell your doctor right away if you get any of these symptoms of PML:
 - weakness on one side of the body that gets worse
 - clumsiness in your arms or legs
 - vision problems
 - changes in thinking and memory
 - confusion
 - personality changes
- herpes zoster infections (shingles), including central nervous system infections
- other serious infections
- decreases in your white blood cell count. Your doctor should do a blood test to check your white blood cell count before you start treatment with BAFIERTAM and while you are on therapy. You should have blood tests after 6 months of treatment and every 6 to 12 months after that.
- **liver problems.** BAFIERTAM may cause serious liver problems that may lead to liver failure, a liver transplant, or death. Your doctor should do blood tests to check your liver function before you start taking BAFIERTAM and during treatment if needed.
 - Tell your doctor right away if you get any of these symptoms of a liver problem during treatment:
 - severe tiredness
 - loss of appetite
 - pain on the right side of your stomach
 - have dark or brown (tea color) urine
 - yellowing of your skin or the white part of your eyes



- **serious gastrointestinal problems,** including bleeding, ulcers, blockage, and tears (perforation) of the stomach or intestines.
 - Tell your healthcare provider right away if you have any of these symptoms during treatment:
 - Stomach-area pain that does not go away
 - Bright red or black stools (that look like tar)
 - Severe vomiting
 - Severe diarrhea
 - Coughing up blood or blood clots
 - Vomiting blood or your vomit looks like "coffee grounds"

The most common side effects of BAFIERTAM[®] include:

- flushing, redness, itching, or rash
- nausea, vomiting, diarrhea, stomach pain, or indigestion
- Flushing and stomach problems are the most common reactions, especially at the start of treatment, and may decrease over time. Call your doctor if you have any of these symptoms and they bother you or do not go away. Ask your doctor if taking aspirin before taking BAFIERTAM may reduce flushing.

These are not all the possible side effects of BAFIERTAM. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

For more information go to dailymed.nlm.nih.gov.

This information does not take the place of talking with your doctor about your medical condition or your treatment.



"For my active lifestyle, BAFIERTAM works. **It gives me the confidence** that I can do the things I like to do." – Annette, living with RMS since 1998



See Why Patients Like You Prefer BAFIERTAM BAFIERTAM.com/patient-stories



You deserve an easy, reliable way to treat MS. Think BAFIERTAM, Pure & Simple

- Small, easy-to-take gel capsules
- No complex tests or exams required to get started
- No dietary or alcohol restrictions
- No known drug interactions
- Easy on, easy off
- Comprehensive patient support

Contact your doctor today and discover if BAFIERTAM is right for you

Banner Life Sciences[®] is the maker of BAFIERTAM. We are here to help you live better with MS, whether you need support navigating insurance, or learning more about your medication.



©2024 Banner. All rights reserved. BAFI-EN-0824-PAT-01D 08/24 BAFIERTAM[®] and the Banner logo are trademarks of Banner Life Sciences[®]. All other trademarks are the property of their respective owners.